

Fall 10-16-2012

Impaired Spatial Learning in Otoconia-Deficient Mice

Ryan M. Yoder

Indiana University - Purdue University Fort Wayne, yoderRM@ipfw.edu

Seth L. Kirby

Indiana University - Purdue University Fort Wayne

Follow this and additional works at: http://opus.ipfw.edu/psych_facpres



Part of the [Psychology Commons](#)

Opus Citation

Ryan M. Yoder and Seth L. Kirby (2012). *Impaired Spatial Learning in Otoconia-Deficient Mice*. Presented at Neuroscience 2012, New Orleans, LA.

http://opus.ipfw.edu/psych_facpres/105

This Poster Session is brought to you for free and open access by the Department of Psychology at Opus: Research & Creativity at IPFW. It has been accepted for inclusion in Psychology Faculty Presentations by an authorized administrator of Opus: Research & Creativity at IPFW. For more information, please contact admin@lib.ipfw.edu.

Introduction

- Damage to head direction cell-containing brain structures causes deficits in spatial learning and navigation, suggesting the head direction signal is important for spatial performance.¹
- Otoconia-deficient *tilted* mice have intact brains with degraded head direction cell signals.²

Aim

To determine whether the degraded head direction signal of *tilted* mice is associated with spatial deficits.


Methods

Subjects

- Male homozygous *tilted* mice and heterozygous littermate controls

Apparatus

- 6-arm Radial Maze in open room



Procedure

1. Pre-exploration:


- All arms baited; maze in a different room
- One 10-min trial per day, for two days

2. Training:

- Two arms baited
- Four trials per day, for ten days

Results

Percentage of Correct Arm Choices



Group:

$$F(1,12) = 64.8, p < .01$$


Trial Block:

$$F(9,108) = 14.2, p < .01$$

Group x Trial Block:

$$F(9,108) = 5.06, p < .01$$

Reference Memory Errors



Group:

$$F(1,12) = 38.9, p < .01$$


Trial Block:

$$F(9,108) = 8.80, p < .01$$

Group x Trial Block:

$$F(9,108) = 7.02, p < .01$$

Working Memory - Correct



Group:

$$F(1,12) = 11.1, p < .01$$


Trial Block (n.s.):

$$F(9,108) = 1.07, p = .39$$

Group x Trial Block:

$$F(9,108) = 2.68, p < .01$$

Working Memory - Incorrect



Group:

$$F(1,12) = 5.73, p = .03$$

Trial Block:

$$F(9,108) = 2.01, p = .04$$

Group x Trial Block:

$$F(9,108) = 4.78, p < .01$$

Summary

Reference memory, working memory-correct, and working memory-incorrect errors occurred more frequently in *tilted* mice than in control mice.

Conclusion

Head direction signal degradation is associated with spatial deficits.

References

1. Taube JS, Kesslak JP, Cotman CW (1992) Lesions of the rat postsubiculum impair performance on spatial tasks. *Behavioral and Neural Biology* 57: 131-143
- Aggleton JP, Hunt PR, Nagle S, Neave N (1996) The effects of selective lesions within the anterior thalamic nuclei on spatial memory in the rat. *Behavioural Brain Research* 81: 189-198.
- Reviews:
 - Muir GM, Taube JS (2002) The neural correlates of navigation: Do head direction cells and place cells guide spatial behavior? *Behavioral and Cognitive Neuroscience Reviews* 1: 297-313;
 - Taube JS (2007) The head direction signal: Origins and sensory-motor integration. *Annual Review of Neuroscience* 30: 181-207
 - Yoder RM, Clark BJ, Taube JS (2011) Origins of landmark encoding in the brain. *Trends in Neurosciences* 34: 561-571.
2. Yoder RM, Taube JS (2009) Head direction cell activity in mice: Robust directional signal depends on intact otolith organs. *Journal of Neuroscience* 29: 1061-1076.

Funding

- NIDCD R15 DC012630
- IPFW Faculty Summer Grant